

DOSAGE & ADMINISTRATION

Clinical studies demonstrated that NOCDURNA was effective at these low doses, specially designed for men and women.¹

Take 1 tablet
daily, 1 hour
before bedtime,
without water



MEN
55.3 mcg/day



WOMEN
27.7 mcg/day



Packaging is not actual size.

WHEN TAKING NOCDURNA

Keep under
tongue until
fully dissolved¹

Limit fluids, starting
from 1 hour before
taking NOCDURNA
and for 8 hours
after taking it¹

Avoid caffeine
or alcohol
before bedtime¹

Empty bladder
immediately
before bedtime¹

INDICATION

NOCDURNA is a vasopressin analog indicated for the treatment of nocturia due to nocturnal polyuria in adults who awaken at least 2 times per night to void.

IMPORTANT SAFETY INFORMATION, INCLUDING BOXED WARNING

WARNING: HYPONATREMIA

See full Prescribing Information for complete boxed warning

- NOCDURNA can cause hyponatremia, which may be life-threatening if severe.
- NOCDURNA is contraindicated in patients at increased risk of severe hyponatremia, such as patients with excessive fluid intake, illnesses that can cause fluid or electrolyte imbalances, and in those using loop diuretics or systemic or inhaled glucocorticoids.
- Ensure serum sodium concentration is normal before starting or resuming NOCDURNA. Measure serum sodium within 1 week and approximately 1 month after initiating therapy and periodically during treatment. More frequently monitor serum sodium in patients 65 years of age and older and in patients at increased risk of hyponatremia.
- If hyponatremia occurs, NOCDURNA may need to be temporarily or permanently discontinued.

Please see additional Important Safety Information on next page.

Click for full Prescribing Information, including complete BOXED WARNING.


SERUM SODIUM MONITORING

Ensure sodium concentration is normal prior to initiating or resuming NOCDURNA. NOCDURNA is contraindicated in patients with hyponatremia or a history of hyponatremia.¹

CHECK SODIUM LEVELS AT THESE TIMEPOINTS¹



BEFORE
STARTING
OR RESUMING
THERAPY



1 WEEK
WITHIN 1 WEEK
OF STARTING
OR RESUMING
THERAPY



1 MONTH
WITHIN 1
MONTH OF
STARTING OR
RESUMING
THERAPY



PERIODICALLY
AS CLINICALLY
APPROPRIATE*

*Patients 65 and older and those at increased risk for hyponatremia should be monitored more frequently.¹

NOCDURNA may need to be temporarily or permanently discontinued if the patient develops hyponatremia, and treatment for hyponatremia instituted, depending on the clinical circumstances, including the duration and severity of the hyponatremia.¹

IMPORTANT SAFETY INFORMATION (cont)

CONTRAINDICATIONS

- Hyponatremia or a history of hyponatremia
- Polydipsia
- Concomitant use with loop diuretics or systemic or inhaled glucocorticoids
- Estimated glomerular filtration rate below 50 mL/min/1.73 m²
- Syndrome of inappropriate antidiuretic hormone secretion (SIADH)
- During illnesses that can cause fluid or electrolyte imbalance
- Heart failure
- Uncontrolled hypertension

WARNINGS AND PRECAUTIONS

- Limit fluid intake to a minimum from 1 hour before until 8 hours after administration. Treatment without concomitant reduction of fluid intake may lead to fluid retention and hyponatremia.
- Fluid retention: Not recommended in patients at risk of increased intracranial pressure or history of urinary retention.

ADVERSE REACTIONS

Common adverse reactions (>2% incidence) included dry mouth, hyponatremia or blood sodium decreased, and dizziness.

DRUG INTERACTIONS

Monitor serum sodium more frequently when NOCDURNA is concomitantly used with drugs that may increase the risk of hyponatremia (e.g. tricyclic antidepressants, selective serotonin re-uptake inhibitors, chlorpromazine, opiate analgesics, thiazide diuretics, carbamazepine, lamotrigine, sulfonyleureas, particularly chlorpropamide and NSAIDs).

USE IN SPECIFIC POPULATIONS

Pregnancy: Use of NOCDURNA is not recommended.

Geriatric Use: Increased risk of hyponatremia if 65 years of age or older. Monitor serum sodium more frequently.

To report SUSPECTED ADVERSE REACTIONS, contact Ferring at 1-888-337-7464 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

Click for full Prescribing Information, including complete BOXED WARNING.

Reference: 1. NOCDURNA [package insert]. Parsippany, NJ: Ferring Pharmaceuticals Inc.